



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A SUSPECTED NOVA IN N. G. C. 2841

A photograph, with an exposure of two hours, made at the 25-foot focus of the 60-inch reflector on February 19, 1912, shows a relatively bright nebulous star in the spiral nebula N. G. C. 2841 that does not appear on earlier and later plates made at the Lick Observatory. The object is approximately 50" west and 20" north of the nucleus of the spiral; it is shown distinctly in the reproduction in the *Astrophysical Journal*, 46, Plate V a, 1917 (*Mt. Wilson Contr.*, No. 132). Professor Curtis independently found the star from an examination of the published photographs. He writes that on plates made with the Crossley reflector on April 17, 1901 (reproduced in *Lick Publication*, 8, Plate 19), and November 19, 1914, no trace of the star is seen. Observations will be continued when the spiral comes into position again.

FRANCIS G. PEASE.

A FAINT NOVA IN THE ANDROMEDA NEBULA.

Two photographs made with the 60-inch reflector on Sept. 11, 1917, show a new star of magnitude 17.5 at a distance of 10' from the center of the Andromeda Nebula (N. G. C. 224.) The approximate position with respect to the nucleus of the nebula is

$$\Delta\alpha = +360''; \quad \Delta\delta = +480''$$

An eighteenth magnitude companion is south 5" and west 2".

The star does not appear on a plate made by Mr. Pease August 21, 1917, and at that time must have been fainter than magnitude 19. No trace of it is found on any of the plates of longer exposure listed by Mr. Ritchey in a preceding note. A photograph kindly made by Mr. Ritchey on Sept. 17, 1917, for the purpose of confirmation, shows the star to have decreased about half a magnitude in six days.

HARLOW SHAPLEY.

NOTE ON THE MAGNITUDES OF NOVAE IN SPIRAL NEBULAE.

Since Ritchey's discovery of the nova in N. G. C. 6946, the number of temporary stars known to have occurred in nebulae has increased so rapidly that an opportunity is afforded of making preliminary use of relative magnitudes in estimating the order of the distances of spiral systems. The results can hope only to be suggestive, but any data will be welcome that bear directly on